Table 5-10. Summary statistics of annual average flow and annual average baseflows, 1933-1942, for USGS Gauge 0224600, NORTH FORK BLACK CREEK NEAR MIDDLEBURG, FL

Statistic	Observed Annual Average Flow (cfs)	HYSEP Local Minimum Annual Average Baseflow (cfs)	BFI Standard Annual Average Baseflow (cfs)	BFI Modified Annual Average Baseflow (cfs)	USF Method Annual Average Baseflow (cfs)
25 th Percentile	181	61	42	42	17
Median	131	50	36	36	14
75 th Percentile	109	43	30	29	12
Minimum	93	37	29	28	8
Maximum	221	77	59	59	19

Table 5-11. Summary statistics of annual average flow and annual average baseflows, 1933-1942, for USGS Gauge 02315500, SUWANNEE RIVER AT WHITE SPRINGS, FL

Statistic	Observed Annual Av- erage Flow (cfs)	HYSEP Local Minimum Annual Average Baseflow (cfs)	BFI Standard Annual Average Baseflow (cfs)	BFI Modified Annual Average Baseflow (cfs)	USF Method Annual Average Baseflow (cfs)
25 th Percentile	1,620	889	1,112	1,123	192
Median	775	469	520	507	106
75 th Percentile	566	320	322	312	57
Minimum	218	99	93	90	22
Maximum	2,283	1,123	1,299	1,370	390

Table 5-12. Summary statistics of annual average flow and annual average baseflows, 1933 -1942, for USGS Gauge 02319000, WITHLACOOCHEE RIVER NEAR PINETTA, FL

Statistic	Observed Annual Average Flow (cfs)	HYSEP Local Minimum Annual Average Baseflow (cfs)	BFI Standard Annual Average Baseflow (cfs)	BFI Modified Annual Average Baseflow (cfs)	USF Method Annual Average Baseflow (cfs)
25 th Percentile	1,759	907	932	932	241
Median	1,129	644	512	514	197
75 th Percentile	577	400	368	380	145
Minimum	413	225	250	221	109
Maximum	2,212	1,158	1,213	1,215	317